

Comment: To prepare for the final exam, you should study these problems, the homework from chapters 26 - 28, as well as reviewing the material from quizzes 1 - 4. The final exam will be comprehensive, but with an emphasis on untested material.

1. Chapter 26, exercise set F, problems 4 and 5.

See solutions at the back of the book.

2. Every day, the quality control engineer for ACME Dairies randomly selects 25 half-gallon (64 fl oz) cartons of whole milk from the day's production run and carefully measures the quantity of milk in each one. If the average amount of milk in this sample differs *significantly* from 64 fl oz, at the 1% significance level, she recalibrates the carton-filling apparatus.

- (a) What does '*...significantly... at the 1% significance level*' mean?

This is statistical significance. In this case, it means that assuming the null hypothesis, the probability is 1% or less, that the difference between the sample average and the (null-hypothetical) expected average is as or more extreme than the observed difference.

- (b) State the null and alternative hypotheses for this test in terms of the appropriate parameter.

H_0 : *The average amount of milk in the daily run of half-gallon cartons is 64 ounces.*

H_1 : *The average amount of milk in the daily run of half-gallon cartons is **not** 64 ounces.*

*(A **two-tailed** test — H_1 does not specify whether the average is greater than or less than 64, just that it is different than 64.)*

- (c) Today's sample of 25 cartons has an average of 64.21 fl oz with a standard deviation of 0.37 fl oz. What is the test statistic? What is the p-value?

The test statistic here is

$$t^* = \frac{(\text{observed average}) - (H_0\text{-expected average})}{SE(\text{average})} = \frac{64.21 - 64}{SD^+/\sqrt{25}} \approx \frac{0.21}{0.378/5} \approx 2.78$$

Comments: *The sample size $n = 25$ is relatively small and we do not know the SD of the error-box, so we approximate it with the SD^+ of the sample, this also implies that the test statistic follows the t -distribution with $25 - 1 = 24$ degrees of freedom.*

The p-value is estimated by looking in the row for 24 degrees of freedom in the t -table in the textbook. The area to the right of $t^ = 2.78$ is between 1% and 0.5% (because $2.49 < 2.78 < 2.80$). This is a **two-tailed** test, because we don't have an expected direction for the difference — a priori, it could be positive or negative. This means that the p-value is two times the entry in the table, i.e.,*

$$1\% < p^* < 2\%.$$

(Using more precise tools, you would find that $p^ \approx 1.04\%$.)*

- (d) What do you conclude?

Since $p^ > 1\%$ (though it's very close!), the ACME Dairies protocol says that the machines do **not** need to be recalibrated.*

- (e) What additional assumptions, if any, are needed to justify the methods (and conclusions) of this test of significance?

To justify the use of the t -test, we must assume that the error-box associated with the filling of the milk cartons follows the normal curve (once the errors are converted to standard units).

3. A marketing firm wants to survey a sample of 1400 adults from a large state to estimate the percentage of adults in the state who prefer streaming movies to watching them on cable TV. Are they more likely to use (a) a *simple random sample*, (b) a *convenience sample* or (c) a *multistage cluster sample* to choose the sample?

Explain your answer.

Ideally, they would like to use a simple random sample, for which the standard errors are smallest, but these are difficult to produce in practice from large populations. From a practical point of view of the three options given, they are most likely to use a multistage cluster sample (if they care about accuracy).

4. A researcher claims to have found a strong correlation ($r = 0.88$) between a person's blood alcohol content (BAC), one hour after drinking, and the type of alcohol they consume (beer, wine or hard liquor).

What is wrong with the researcher's claim? What would make more sense here? Explain.

The correlation does not make sense because you can't compute the correlation between BAC (numerical) and alcohol type (nonnumerical). There is no way to interpret the number 0.88 (and it is not clear how it was computed).

*The simplest correlation that would make sense here is a correlation between the amount of alcohol consumed and the BAC an hour later. If one wants to study the effect of the **type** of alcoholic beverage and BAC an hour later, a more involved study is needed.*

*E.g., the researcher could study the effect of each type of alcoholic beverage separately and **control for various confounding factors**, like the amount of alcohol that is consumed, weight and gender of the drinker, etc.*

5. Investigators studying the relationship between cigarette smoking and blood pressure in adult men collected data from 6235 U.S. men aged 20 - 40, and generated the following statistics:

$$\begin{aligned} \bar{X} &= 24 & SD_X &= 5.5 \\ \bar{Y} &= 135 & SD_Y &= 9 & r &= 0.7 \end{aligned}$$

where X = number of cigarettes per day, and Y = systolic blood pressure, measured in mmHG.

- (a) Use the *regression method* to estimate the average systolic blood pressure for U.S. men, aged 20 - 40 who smoke 20 cigarettes per day. *Show your work.*

*20 cigarettes per day is $(24 - 20)/5.5 \approx 0.727 SD_x$ **below** average, so the regression method predicts that the average systolic blood pressure of men who smoke 20 cigarettes per day will be*

$$0.727 \times r \times SD_y = 0.727 \times 0.7 \times 9 \approx 4.58 \text{ mmHG}$$

***below** the average of 135 mmHG. I.e., the predicted average blood pressure for these men is about 130.42 mmHG.*

- (b) What is the predicted systolic blood pressure of a 28-year old man who smokes 30 cigarettes per day? Include a 'give-or-take' number with your estimate. *Show your work.*

*30 cigarettes per day is $(30 - 24)/5.5 \approx 1.09 SD_x$ **above** average. The regression method says that the average blood pressure of men (aged 20 - 40) who smoke this much is predicted to be*

$$1.09 \times 0.7 \times 9 \approx 6.87 \text{ mmHG}$$

***above** average, or about 141.87 mmHG.*

The SER (root mean square error of regression) for predicting blood pressure from cigarettes per day is

$$SER = \sqrt{1 - 0.7^2} \times SD_y \approx 6.43,$$

so the blood pressure of an **individual** 28 year old man who smokes 30 cigarettes per day is predicted to be about 141.87 ± 6.43 mmHG.

- (c) Joseph is a 60-year old man who smokes about 40 cigarettes a day. Is it reasonable to predict that his systolic blood pressure is somewhere between 147 and 160 mmHG, based on the given information? *Explain your answer.*

This is not a question of whether the calculations were done correctly. The data was collected from men aged 20 - 40, and cannot be used to predict blood pressure for a man whose age is so far outside the range of ages in the study. Especially given the possible effects of age on blood pressure (blood pressure tends to increase with age).

6. John Smith is running for office. One week before the election, his campaign manager hires a Polling firm to survey likely voters. The firm surveyed a simple random sample of 2700 likely voters and found that 51% favor Smith. They also found that of the 1250 women in the survey, 54% favor Smith.

You may assume that the survey was based on a simple random sample, that the population is in the millions and that to win the office, the candidate needs to win more than 50% of the votes cast.

- (a) What percentage of the men in the survey favor Smith?

675 = 54% × 1250 of the women surveyed favored Smith, and a total of 51% × 2700 = 1377 of the people surveyed favored him. So, 1377 - 675 = 702 men surveyed favored Smith. A total of 2700 - 1250 = 1450 men were surveyed, so

$$\frac{702}{1450} \times 100\% \approx 48.41\%$$

of the men surveyed favor Smith.

- (b) Compute 95% confidence intervals for the percentage of women who favor Smith, the percentage of men who favor Smith and the percentage of likely voters who favor Smith.

Women: *The observed percentage is 54%, and the standard error is*

$$SE_W = \frac{\sqrt{0.54 \times 0.46}}{\sqrt{1250}} \times 100\% \approx 1.41\%.$$

The 95% confidence interval for the percentage of women who favor Smith is

$$(54\% \pm 2SE_W) = (54\% \pm 2.82\%).$$

Men: *The observed percentage is 48.41%, and the standard error is*

$$SE_M = \frac{\sqrt{0.4841 \times 0.5159}}{\sqrt{1450}} \times 100\% \approx 1.31\%.$$

The 95% confidence interval for the percentage of men who favor Smith is

$$(48.41\% \pm 2SE_M) = (48.41\% \pm 2.62\%).$$

All: *The observed percentage is 51%, and the standard error is*

$$SE_A = \frac{\sqrt{0.51 \times 0.49}}{\sqrt{2700}} \times 100\% \approx 0.96\%.$$

The 95% confidence interval for the percentage of all likely voters who favor Smith is

$$(51\% \pm 2SE_A) = (51\% \pm 1.92\%).$$

7. As part of a class project, a statistics student at a large university (15,000 students — 9000 men and 6000 women), went to the central plaza of the campus at noon one day, approached 100 students and asked them where they went to high school. His sample included 51 women and 49 men. Is it likely that the student's sampling procedure was like taking a simple random sample? Justify your answer as precisely as possible (using numbers, probability, etc.).

If the student's sampling procedure was like taking a simple random sample, then it was like drawing 100 tickets at random, without replacement from a 0-1 box of 15000 tickets, where 60% of the tickets are $\boxed{1}$ s and 40% are $\boxed{0}$ s. The question now becomes:

How likely is it to draw 49% $\boxed{1}$ s (or less) from a box with 60% $\boxed{1}$ s, in 100 random draws?

To answer this question, we use the **Normal Approximation**. The SD of the box is

$$SD = \sqrt{0.6 \times 0.4} \approx 0.49,$$

and the $SE_{\%}$ for 100 draws from this box is

$$SE_{\%} = \frac{0.49}{\sqrt{100}} \cdot 100\% = 4.9\%.$$

(Technically, the $SE_{\%}$ is **slightly** smaller, because the draws are done without replacement, but since there are 15000 tickets in the box and only 100 are drawn, the correction factor is very close to 1.)

According to the normal approximation, the probability of drawing 49% (or fewer) $\boxed{1}$ s from this box with a simple random sample is about equal to the area under the normal curve to the left of

$$z = \frac{49\% - 60\%}{4.9\%} \approx -2.24,$$

which is approximately 1.25%.

To summarize, the probability that a simple random sample of students from this University would have 49% men (or fewer) is about 1.25%, and we can conclude that the student's sample in this case was almost certainly **not** a simple random sample.

Indeed, from the description, it is clear that this was a **sample of convenience** (and perhaps biased towards women).

8. According to the 1999 census, the median household income in the city of San Diego was \$46,500. In 2004, a high-end grocery chain hires a statistical research firm to corroborate their marketing consultant's claim that median household income has gone up since 1999. The research firm takes a simple random sample of 600 San Diego households and finds that 55% of the sample households have incomes above \$46,500.

Was the consultant right? Frame your answer in terms of an appropriate test of significance.

To answer the question, we use a test of significance.

- **Null hypothesis:** The median income has not gone up since 1999. I.e., 50% of the households in San Diego have incomes above \$46,500, (and 50% have incomes below this level).

Alternative hypothesis: The median income has gone up, so more than 50% of the households have incomes above \$46,500.

- **Null-hypothetical box model:** A 0-1 box with a $\boxed{1}$ for every household in San Diego (in 2004) with income above \$46,500 and a $\boxed{0}$ for every household in San Diego (in 2004) with income below \$46,500. The null hypothesis says that 50% of the tickets in this box are $\boxed{1}$ s.

- **Data:** The sample percentage of households with incomes above \$46,500 is 55%.
- **Test statistic:** The observed percentage is 55% and the null-hypothetical expected percentage is 50%. Furthermore, the SD of the null-hypothetical box is $\sqrt{1/2 \times 1/2} = 1/2$, so the standard error is $SE_{\%} = \frac{0.5}{\sqrt{600}} \times 100\% \approx 2.04\%$.

Hence the test statistic is

$$z = \frac{\text{observed \%} - \text{expected \%}}{SE_{\%}} = \frac{55\% - 50\%}{2.04\%} \approx 2.45.$$

- **P-value** (observed significance level): The P-value here is the area under the normal curve to the right of $z = 2.45$ which is about $\frac{100\% - 98.57\%}{2} \approx 0.715\%$.
- **Conclusion:** The P-value is very low (less than 1%), so we reject the null hypothesis and conclude that the consultant was right — the median income in 2004 was higher than \$46,500.

9. Suppose that a fair die is rolled 3 times.

a. What is the probability that a \square is observed **at least once**?

The probability of no \square s in 3 rolls is

$$\frac{5}{6} \cdot \frac{5}{6} \cdot \frac{5}{6} \approx 57.87\%$$

so the probability of **at least one** \square in three rolls is $100\% - 57.87\% = 42.13\%$.

b. What is the probability that a \square is observed **exactly once**?

If a \square is observed exactly once, then...

... it occurs on roll one, but not on rolls two and three; or it occurs on roll two, but not on rolls one or three; or it occurs on roll three, but not on rolls one or two.

Each of these three possibilities has the same probability, namely

$$\frac{1}{6} \cdot \frac{5}{6} \cdot \frac{5}{6} \approx 0.11574,$$

and all three are **mutually exclusive** so the probability that exactly one of them occurs is

$$\frac{1}{6} \cdot \frac{5}{6} \cdot \frac{5}{6} + \frac{1}{6} \cdot \frac{5}{6} \cdot \frac{5}{6} + \frac{1}{6} \cdot \frac{5}{6} \cdot \frac{5}{6} \approx 0.3472.$$

c. What is the probability of that the **sum** of the three rolls is 4 or 5?

First, we need to find the different configurations of three dice that result in sums of 4 or 5.

(i) The only way to obtain a sum of 4 in three rolls is with one \square and two \square s. This can occur in three ways: $\square\square\square$, $\square\square\square$ or $\square\square\square$. Each of these configurations has the same probability, namely $\frac{1}{6} \cdot \frac{1}{6} \cdot \frac{1}{6} = 1/216$, and they are all mutually exclusive, so the probability of a sum of 4 in three rolls is

$$3 \times \frac{1}{216} = \frac{1}{72} \approx 1.389\%.$$

(ii) The only ways to obtain a sum of 5 in three rolls is with (a) two \square s and one \square or (b) one \square and two \square s. In other words, the only way to obtain a sum of 5 is with one of the configurations

$$\square\square\square, \square\square\square, \square\square\square, \square\square\square, \square\square\square \text{ or } \square\square\square.$$

Each of these six configurations has the same probability, namely $\frac{1}{6} \cdot \frac{1}{6} \cdot \frac{1}{6} = 1/216$, and they are all mutually exclusive so the probability of a sum of 5 in three rolls is

$$6 \times \frac{1}{216} = \frac{1}{36} \approx 2.778\%.$$

Finally, since a sum of 4 and a sum of 5 are mutually exclusive, the probability of a sum of 4 or a sum of 5 in three rolls is

$$\frac{1}{72} + \frac{1}{36} = \frac{3}{72} = \frac{1}{24} \approx 4.167\%.$$

10. Suppose that a fair die is rolled 600 times.

a. What is the expected number of $\square\bullet$ s?

Given that the die is fair, the probability of observing a $\square\bullet$ on any given roll is $1/6$, and the **expected number** of $\square\bullet$ s is therefore equal to

$$\frac{1}{6} \cdot 600 = 100.$$

b. What is the probability that a $\square\bullet$ is observed between 95 and 105 times?

The SD of the 'die-box' is $\sqrt{1/6 \times 5/6} \approx 0.373$, and the SE for the **number** of $\square\bullet$ s in 600 draws is $SD \times \sqrt{600} \approx 9.129$. By the normal approximation, the probability of observing between 95 and 105 $\square\bullet$ s in 600 draws is therefore approximately equal to the area under the normal curve between

$$\frac{94.5 - 100}{9.129} \approx -0.60 \quad \text{and} \quad \frac{105.5 - 100}{9.129} \approx 0.60$$

which is 45.15%.

c. What is the probability that more than 110 $\square\bullet$ s are observed?

Once again, we invoke the normal approximation and conclude that this probability is approximately equal to the area under the normal curve to the right of

$$\frac{110.5 - 100}{9.129} \approx 1.15$$

which is

$$\frac{100\% - 74.99\%}{2} = 12.505\%.$$

d. What is the probability that the **sum** of the 600 rolls is between 2070 and 2130?

Once again, we invoke the normal approximation, but this time it is for the sum of draws from the box $\boxed{1 \ 2 \ 3 \ 4 \ 5 \ 6}$.

The average of this box is $(1 + 2 + 3 + 4 + 5 + 6)/6 = 3.5$ and the SD of this box is

$$\sqrt{\frac{(1 - 3.5)^2 + (2 - 3.5)^2 + (3 - 3.5)^2 + (4 - 3.5)^2 + (5 - 3.5)^2 + (6 - 3.5)^2}{6}} \approx 1.708.$$

The expected value of the sum of 600 draws from this box is $600 \times 3.5 = 2100$ and the standard error for the sum of 600 draws is $SE = \sqrt{600} \times SD \approx 41.833$. Using the normal approximation, we find that

$$P(2070 \leq \text{sum of 600 rolls} \leq 2130) \approx \text{area under NC between } \frac{2070 - 2100}{41.833} \text{ and } \frac{2130 - 2100}{41.833}.$$

Now, $\frac{2070 - 2100}{41.833} \approx -0.717$ and $\frac{2130 - 2100}{41.833} \approx 0.717$, so the probability we seek is about 53% (between the table entries for 0.70 and 0.75, closer to the one for 0.70 and rounded).

11. There are about 25,000 high schools in the United States and each high school has a principal. These 25,000 high schools also employ a total of about one million teachers. As part of a national survey of education, a simple random sample of 625 high schools is chosen.

- (a) In 505 of the sample high schools the principal has an advanced degree. If possible, find an approximate 95% confidence interval for the percentage of all 25,000 high school principals who have advanced degrees. If this is not possible, explain why not.

The sample percentage of principals with advanced degrees is $505/625 \times 100\% = 80.8\%$. The sample SD in this case is $\sqrt{0.808 \times 0.192}$, so the Standard error (for percentage) is

$$SE_{\%} \approx \frac{\sqrt{0.808 \times 0.192}}{\sqrt{625}} \times 100\% \approx 1.58\%.$$

Hence a 95% confidence interval for the percentage of high schools whose principal has an advanced degree is

$$80.8\% \pm 2 \times 1.58\% = 80.8\% \pm 3.16\% \quad \text{or } (76.64\%, 83.96\%).$$

- (b) As it turned out, the 625 sample high schools described above employed a total of 12,000 teachers, of whom 6,500 had advanced degrees. If possible, find an approximate 95% confidence interval for the percentage of all one million high school teachers with advanced degrees. If this is not possible, explain why not.

*The sample of 12,000 teachers in this hypothetical example is **not** a simple random sample of U.S. high school teachers — taking all of the teachers from a random sample of high schools is not the same thing as a random sample of teachers from the whole country — it is a cluster sample of teachers. The methods we have been using do not apply in this case, and we cannot find a 95% confidence interval using these methods.*

12. A researcher studying the media consumption habits of U.S. adults suspects that women watch more ‘reality’ shows than men. To test this hypothesis, she surveys a simple random sample of 1225 U.S. men and a simple random sample of 1444 U.S. women. The men surveyed watched an average of 4.36 hours per week of ‘reality’ shows, with an SD of 1.8 hours per week. The women watched an average of 4.43 hours per week of ‘reality’ shows, with an SD of 1.7 hours per week.

- (a) Formulate appropriate null and alternative hypotheses in terms of a box model to test the researcher’s hypothesis at the 5% significance level.

The null hypothesis says that the women and men watch the same amount of reality shows, on average. I.e., if μ_w is the average number of hours/week that women watch reality shows and μ_m is the average number of hours/week that men watch reality shows, then the researcher’s hypotheses are..

$$H_0 : \mu_w = \mu_m, \text{ and}$$

$$H_1 : \mu_w > \mu_m.$$

The box model is that the box for women and the box for men have the same averages.

- (b) Find the test statistic and the P -value.

The test statistic is

$$z^* = \frac{\bar{w} - \bar{m}}{\sqrt{SE_w^2 + SE_m^2}},$$

where \bar{w} and \bar{m} are the observed averages for women and men respectively, and SE_w and SE_m are the standard errors for women and men respectively. Plugging in the given sample statistics, we find that

$$\bar{w} = 4.43, SE_w = \frac{1.7}{\sqrt{1444}} \approx 0.0447, \bar{m} = 4.36 \text{ and } SE_m = \frac{1.8}{\sqrt{1225}} \approx 0.0514,$$

so

$$z^* = \frac{4.43 - 4.36}{\sqrt{(0.0447)^2 + (0.0514)^2}} \approx 1.03.$$

The p -value is equal to the area under the normal curve to the right of $z^* = 1.03$, which is about 15%.

- (c) Is the researcher right? In what sense? Explain.

The data does not support the researcher's claim. The difference between the averages for men and women is small and as the p -value shows in this case, it can be explained reasonably by chance error.

13. Chapter 27, Review problem 7.

Comment: This problem is similar to the radiation-surgery example in section 4 of chapter 27.

- (a) The observed difference between the rates of recidivism is (control - treatment) $49.4\% - 48.3\% = 1.1\%$. The $SE_{\%}$ for the control group is

$$SE_c = \frac{\sqrt{0.494 \times 0.506}}{\sqrt{154}} \times 100\% \approx 4\%$$

and the $SE_{\%}$ for the treatment group is

$$SE_t = \frac{\sqrt{0.483 \times 0.517}}{\sqrt{592}} \times 100\% \approx 2\%.$$

so the SE for the difference is

$$SE_{diff} = \sqrt{(0.04)^2 + (0.02)^2} \times 100\% \approx 4.5\%.$$

This means that the test statistic is

$$z^* = \frac{1.1\%}{4.5\%} \approx 0.24,$$

and the p -value (from the table) is about 40%.

Conclusion: The observed difference in recidivism rates can be explained by chance — the income support did not seem to have a benefit.

- (b) The observed difference between the average number of weeks worked (control - treatment) is $24.3 - 16.8 = 7.5$. The SE for the control group is

$$SE_c = \frac{17.3}{\sqrt{154}} \approx 1.394$$

and the SE for the treatment group is

$$SE_t = \frac{15.9}{\sqrt{592}} \approx 0.653$$

so the SE for the difference is

$$SE_{diff} = \sqrt{(1.394)^2 + (0.653)^2} \approx 1.54.$$

The test statistic is

$$z^* = \frac{7.5}{1.54} \approx 4.87$$

so the p -value is effectively 0%.

Conclusion: The observed difference in weeks-worked between the control and treatment group is **not** due to chance error. The released prisoners who received income support tended to work less than those who did not. Perhaps this explains the failure of the program to reduce recidivism.